

Due: Sunday, 4/6, 4:00 PM
Grace period until Sunday, 4/6, 6:00 PM

Sundry

Before you start writing your final homework submission, state briefly how you worked on it. Who else did you work with? List names and email addresses. (In case of homework party, you can just describe the group.)

1 Probability Potpourri

Note 13
Note 14

Provide brief justification for each part.

- For two events A and B in any probability space, show that $\mathbb{P}[A \setminus B] \geq \mathbb{P}[A] - \mathbb{P}[B]$.
- Suppose $\mathbb{P}[D | C] = \mathbb{P}[D | \bar{C}]$, where \bar{C} is the complement of C . Prove that D is independent of C .
- If A and B are disjoint, does that imply they're independent?

2 Independent Complements

Note 14

Let Ω be a sample space, and let $A, B \subseteq \Omega$ be two independent events.

- Prove or disprove: \bar{A} and \bar{B} must be independent.
- Prove or disprove: A and \bar{B} must be independent.
- Prove or disprove: A and \bar{A} must be independent.
- Prove or disprove: It is possible that $A = B$.

3 Cliques in Random Graphs

Note 13
Note 14

Consider the graph $G = (V, E)$ on n vertices which is generated by the following random process: for each pair of vertices u and v , we flip a fair coin and place an (undirected) edge between u and v if and only if the coin comes up heads.

- What is the size of the sample space?

- (b) A k -clique in a graph is a set S of k vertices which are pairwise adjacent (every pair of vertices is connected by an edge). For example, a 3-clique is a triangle. Let E_S be the event that a set S forms a clique. What is the probability of E_S for a particular set S of k vertices?
- (c) Suppose that $V_1 = \{v_1, \dots, v_\ell\}$ and $V_2 = \{w_1, \dots, w_k\}$ are two arbitrary sets of vertices. What conditions must V_1 and V_2 satisfy in order for E_{V_1} and E_{V_2} to be independent? Prove your answer.
- (d) Prove that $\binom{n}{k} \leq n^k$. (You might find this useful in part (e)).
- (e) Prove that the probability that the graph contains a k -clique, for $k \geq 4\log_2 n + 1$, is at most $1/n$. *Hint:* Use the union bound.

4 Poisoned Smarties

Note 14

Supposed there are 3 people who are all owners of their own Smarties factories. Burr Kelly, being the brightest and most innovative of the owners, produces considerably more Smarties than her competitors and has a commanding 50% of the market share. Yousef See, who inherited her riches, lags behind Burr and produces 40% of the world's Smarties. Finally Stan Furd, brings up the rear with a measly 10%. However, a recent string of Smarties related food poisoning has forced the FDA investigate these factories to find the root of the problem. Through her investigations, the inspector found that 2 Smarties out of every 100 at Kelly's factory was poisonous. At See's factory, 5% of Smarties produced were poisonous. And at Furd's factory, the probability a Smarty was poisonous was 0.1.

- (a) What is the probability that a randomly selected Smarty will be safe to eat?
- (b) If we know that a certain Smarty didn't come from Burr Kelly's factory, what is the probability that this Smarty is poisonous?
- (c) If a randomly selected Smarty is poisonous, what is the probability it came from Stan Furd's Smarties Factory?

5 Symmetric Marbles

Note 14

A bag contains 4 red marbles and 4 blue marbles. Rachel and Brooke play a game where they draw four marbles in total, one by one, uniformly at random, without replacement. Rachel wins if there are more red than blue marbles, and Brooke wins if there are more blue than red marbles. If there are an equal number of marbles, the game is tied.

- (a) Let A_1 be the event that the first marble is red and let A_2 be the event that the second marble is red. Are A_1 and A_2 independent?
- (b) What is the probability that Rachel wins the game?
- (c) Given that Rachel wins the game, what is the probability that all of the marbles were red?

Now, suppose the bag contains 8 red marbles and 4 blue marbles and we add a tiebreaker to the game: if there are an equal number of red and blue marbles among the four drawn, Rachel wins if the third marble is red, and Brooke wins if the third marble is blue.

- (d) What is the probability that the third marble is red?
- (e) Given that there are k red marbles among the four drawn, where $0 \leq k \leq 4$, what is the probability that the third marble is red? Answer in terms of k .
- (f) Given that the third marble is red, what is the probability that Rachel wins the game?

6 Socks

Note 13

Note 14

Suppose you have n different pairs of socks (n left socks and n right socks, for $2n$ individual socks total) in your dresser. You take the socks out of the dresser one by one without looking and lay them out in a row on the floor. In this question, we'll go through the computation of the probability that no two matching socks are next to each other.

- (a) We can consider the sample space as the set of length $2n$ permutations. What is the size of the sample space Ω , and what is the probability of a particular permutation $\omega \in \Omega$?
- (b) Let A_i be the event that the i th pair of matching socks are next to each other. Calculate $\mathbb{P}[A_i]$.
- (c) Calculate $\mathbb{P}[A_1 \cap \dots \cap A_k]$ for an arbitrary $k \geq 2$. (Hint: try using a counting based approach.)
- (d) Putting these all together, calculate the probability that there is at least one pair of matching socks next to each other. Your answer can (and should) be expressed as a summation. (Hint: use Inclusion/Exclusion.)
- (e) Using your answer from the previous part, what is the probability that no two matching socks are next to each other? (This should follow directly from your answer to the previous part, and also can be left as a summation.)