

Due: Saturday 5/3, 4:00 PM
Grace period until Saturday 5/3, 6:00 PM

Sundry

Before you start writing your final homework submission, state briefly how you worked on it. Who else did you work with? List names and email addresses. (In case of homework party, you can just describe the group.)

1 Predictable Gaussians

Note 21 Let Y be the result of a fair coin flip, and X be a normally distributed random variable with parameters dependent on Y . That is, if $Y = 1$, then $X \sim N(\mu_1, \sigma_1^2)$, and if $Y = 0$, then $X \sim N(\mu_0, \sigma_0^2)$.

(a) Sketch the two distributions of X overlaid on the same graph for the following cases:

(i) $\sigma_0^2 = \sigma_1^2, \mu_0 \neq \mu_1$

(ii) $\sigma_0^2 \neq \sigma_1^2, \mu_0 = \mu_1$

(b) Bayes' rule for mixed distributions can be formulated as $\mathbb{P}[Y = 1 | X = x] = \frac{\mathbb{P}[Y=1]f_{X|Y=1}(x)}{f_X(x)}$ where Y is a discrete distribution and X is a continuous distribution. Compute $\mathbb{P}[Y = 1 | X = x]$, and show that this can be expressed in the form of $\frac{1}{1+e^\gamma}$ for some expression γ . (Hint: any value z can be equivalently expressed as $e^{\ln(z)}$)

(c) In the special case where $\sigma_0^2 = \sigma_1^2$ find a simple expression for the value of x where $\mathbb{P}[Y = 1 | X = x] = \mathbb{P}[Y = 0 | X = x] = 1/2$, and interpret what the expression represents. (Hint: the identity $(a+b)(a-b) = a^2 - b^2$ may be useful)

2 Moments of the Gaussian

Note 21 For a random variable X , the quantity $\mathbb{E}[X^k]$ for $k \in \mathbb{N}$ is called the k th moment of the distribution. In this problem, we will calculate the moments of a standard normal distribution.

(a) Prove the identity

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left(-\frac{tx^2}{2}\right) dx = t^{-1/2}$$

for $t > 0$.

Hint: Consider a normal distribution with variance $\frac{1}{t}$ and mean 0.

- (b) For the rest of the problem, X is a standard normal distribution (with mean 0 and variance 1). Use part (a) to compute $\mathbb{E}[X^{2k}]$ for $k \in \mathbb{N}$.

Hint: Try differentiating both sides with respect to t , k times. You may use the fact that we can differentiate under the integral without proof.

- (c) Compute $\mathbb{E}[X^{2k+1}]$ for $k \in \mathbb{N}$.

3 Chebyshev's Inequality vs. Central Limit Theorem

Note 17
Note 21

Let n be a positive integer. Let X_1, X_2, \dots, X_n be i.i.d. random variables with the following distribution:

$$\mathbb{P}[X_i = -1] = \frac{1}{12}; \quad \mathbb{P}[X_i = 1] = \frac{9}{12}; \quad \mathbb{P}[X_i = 2] = \frac{2}{12}.$$

- (a) Calculate the expectations and variances of X_1 , $\sum_{i=1}^n X_i$, $\sum_{i=1}^n (X_i - \mathbb{E}[X_i])$, and


$$Z_n = \frac{\sum_{i=1}^n (X_i - \mathbb{E}[X_i])}{\sqrt{n/2}}.$$

- (b) Use Chebyshev's Inequality to find an upper bound b for $\mathbb{P}[|Z_n| \geq 2]$.
 (c) Use b from the previous part to bound $\mathbb{P}[Z_n \geq 2]$ and $\mathbb{P}[Z_n \leq -2]$.
 (d) As $n \rightarrow \infty$, what is the distribution of Z_n ?
 (e) We know that if $Z \sim \mathcal{N}(0, 1)$, then $\mathbb{P}[|Z| \leq 2] = \Phi(2) - \Phi(-2) \approx 0.9545$. As $n \rightarrow \infty$, provide approximations for $\mathbb{P}[Z_n \geq 2]$ and $\mathbb{P}[Z_n \leq -2]$.

4 Analyze a Markov Chain

Note 22

Consider a Markov chain with the state diagram shown below where $a, b \in (0, 1)$.



Here, we let $X(n)$ denote the state at time n .

- (a) Is this Markov chain irreducible? Is this Markov chain aperiodic? Justify your answers.
 (b) Calculate $\mathbb{P}[X(1) = 1, X(2) = 0, X(3) = 0, X(4) = 1 \mid X(0) = 0]$.
 (c) Calculate the invariant distribution. Do all initial distributions converge to this invariant distribution? Justify your answer.

5 A Bit of Everything

Note 22

Suppose that X_0, X_1, \dots is a Markov chain with finite state space $S = \{1, 2, \dots, n\}$, where $n > 2$, and transition matrix P . Suppose further that

$$P(1, i) = \frac{1}{n} \quad \text{for all states } i \text{ and}$$
$$P(j, j-1) = 1 \quad \text{for all states } j \neq 1,$$

with $P(i, j) = 0$ everywhere else.

- Prove that this Markov chain is irreducible and aperiodic.
- Suppose you start at state 1. What is the distribution of T , where T is the number of transitions until you leave state 1 for the first time?
- Again starting from state 1, what is the expected number of transitions until you reach state n for the first time?
- Again starting from state 1, what is the probability you reach state n before you reach state 2?
- Compute the stationary distribution of this Markov chain.

6 Playing Blackjack

Note 22

Suppose you start with \$1, and at each turn, you win \$1 with probability p , or lose \$1 with probability $1 - p$. You will continually play games of Blackjack until you either lose all your money, or you have a total of n dollars.

- Formulate this problem as a Markov chain.
- Let $\alpha(i)$ denote the probability that you end the game with n dollars, given that you started with i dollars.

Notice that for $0 < i < n$, we can write $\alpha(i+1) - \alpha(i) = k(\alpha(i) - \alpha(i-1))$. Find k .

- Using part (b), find $\alpha(i)$, where $0 \leq i \leq n$. (You will need to split into two cases: $p = \frac{1}{2}$ or $p \neq \frac{1}{2}$.)

Hint: Try to apply part (b) iteratively, and look at a telescoping sum to write $\alpha(i)$ in terms of $\alpha(1)$. The formula for the sum of a finite geometric series may be helpful when looking at the case where $p \neq \frac{1}{2}$:

$$\sum_{k=0}^m a^k = \frac{1 - a^{m+1}}{1 - a}.$$

Lastly, it may help to use the value of $\alpha(n)$ to find $\alpha(1)$ for the last few steps of the calculation.

- As $n \rightarrow \infty$, what happens to the probability of ending the game with n dollars, given that you start with i dollars, with the following values of p ?

- (i) $p > \frac{1}{2}$
- (ii) $p = \frac{1}{2}$
- (iii) $p < \frac{1}{2}$